

Study of prevalence of congenital heart diseases in children

¹Dr. Pankaj kumar Kashyap, ²Dr. Anil Kumar Jaiswal

¹ Senior Resident, Upgraded Department of Paediatrics, Patna Medical College & Hospital, Patna, Bihar, India

² Professor & HOD, Upgraded Department of Paediatrics, Patna medical College & Hospital, Patna, Bihar, India

Abstract

Our aim was to study the prevalence, age-wise distribution, and clinical spectrum of congenital heart disease (CHD) at a P.M.C.H. A retrospective analysis of records of 10,641 patients over a 16 months done. Clinical examination, echocardiography and color doppler were used as diagnostic tools. A prevalence of 26.4 per 1000 patients was observed. VSD (ventricular septal defect) was the commonest lesion (21.3%), followed by ASD (atrial septal defect) in 18.9% and PDA (patent ductus arteriosus) in 14.6%. Tetralogy of Fallot was the commonest cyanotic heart disease (4.6%). Maximum number of children with heart disease (82.9%) were diagnosed between 0-3 years of age.

Keywords: congenital heart disease, India, prevalence

1. Introduction

Congenital heart diseases (CHD) are relatively common with a prevalence ranging from 3.7 to 17.5 per 1000 live births [1, 2]. According to a status report on CHD in India, 10% of the present infant mortality may be accounted for by CHD [3].

According to a large hospital based study from India, the incidence of congenital heart disease is 3.9/ 1000 live births [4]. In community based studies from India [5, 6] the prevalence of CHD ranges from 0.8 – 5.2/1000 patients. Thus, the prevalence of CHD is not uniform across the country and setting. We conducted this study to assess the prevalence of CHD among patients attending a P.M.C.H hospital in Bihar.

2. Methods

This work was done at the Upgraded Department of Pediatrics Patna medical college and hospital at Patna, with patients coming from all over the state with population of 11Crore. Patna has a population of around 2.5 million. We receive patients from lower to upper middle class strata. We retrospectively analyzed the records of all live births in the hospital from march 2015 to June 2016, and all records of pediatric first visit OPD (out patient department) and IPD

(indoor patient department) patients between 0 to 15 years, during the same period.

Children diagnosed with CHD were analyzed further. Pre terms with PDA were followed up and not included in this study if it closed spontaneously within the period of hospital stay or by managing conservatively. Clinical examination, 2D echocardiography and color doppler were considered as definitive tools for diagnosis of CHD.

3. Results

CHD was detected in 9 out of 860 live births in the hospital during the period march 2015 to June 2016. During this period we saw 10,641 new patients between 0 to 15 yrs (OPD and IPD), of these, 281 children were diagnosed to be having some type of CHD, with a prevalence of 26.4 per 1000 patients. Details of congenital heart diseases are provided in Table I. Ventricular septal defect was the most common heart lesion (21.3%). Acyanotic heart diseases were present in 222 (79%) children; 59(21%) had cyanotic heart diseases. Maximum number of cases were seen in 0-3 years age group (n=233, 82.9%).

Table 1: Spectrum of age wise distribution of congenital heart diseases

Age group	0-3yr		4-6yr		7-9yr		10-12 yr		13-15 yr		Total	%
	M	F	M	F	M	F	M	F	M	F		
VSD(subaortic 85%)	23	20	6	2	4	0	2	0	2	1	60	21.3
ASD	36	10	5	0	2	0	0	0	0	0	53	18.9
PDA	22	15	1	1	2	0	0	0	0	0	41	14.6
Complex anomalies	28	5	1	0	0	1	0	0	0	0	35	12.5
AV septal defects	16	12	1	0	0	0	0	0	0	0	29	10.3
ASD+PDA	11	1	1	0	0	0	0	0	0	0	13	4.6
TOF	9	0	1	0	3	0	0	0	0	0	13	4.6
PS	2	2	4	1	0	0	0	0	0	0	9	3.2

AS	2	3	1	0	1	0	0	0	1	0	8	2.8
Ebsteins anomaly	3	0	0	0	0	0	0	0	0	0	3	1.1
TGA+VSD+PS	2	0	0	0	0	1	0	0	0	0	3	1.1
Peripheral PS	3	0	0	0	0	0	0	0	0	0	3	1.1
Bicuspid aortic valve	1	1	1	0	0	0	1	0	0	0	3	1.1
Truncus arteriosus	1	1	0	0	0	0	0	0	0	0	2	0.71
TAPVC	1	0	0	0	0	0	0	0	0	0	1	0.4
Dextroversion	1	0	0	0	0	0	0	0	0	0	1	0.4
Sinus venosus	0	0	1	0	0	0	0	0	0	0	1	0.4
DORV+PS	1	0	0	0	0	0	0	0	0	0	1	0.4
Single ventricle	0	1	0	0	0	0	0	0	0	0	1	0.4
VSD+PDA	1	0	0	0	0	0	0	0	0	0	1	0.4
Total	162	71	23	4	12	2	3	0	3	1	281	

4. Discussion

As a group, congenital heart disease (CHD) constitute a significant proportion (up to 25% in some studies) of congenital malformations that present in the neonatal period. Recent studies from India and other developing countries have shown a decline in prevalence of rheumatic fever and rheumatic heart disease [7].

Congenital malformations and, in particular CHDs are likely to become important contributors to infant mortality in the near future. Hence, it is important to determine the exact prevalence and case burden of congenital heart disease so that appropriate changes in health policies can be recommended [8]. There is just one study available from India which gives the incidence of CHD per 1000 live births, by Khalil, *et al.* [4]. They studied 10964 live births and observed the incidence of 3.9/1000 live births. All large studies from India have taken into consideration only one age group i.e., either newborns or school going children. The former may miss out on a large number of small VSD or Tetralogy of Fallot or ductus dependent lesions, which present a little later than at birth. It also fails to focus on the prevalence of CHD.

Other studies done on the prevalence of CHD in community include mainly 5-15 year old school going children, which automatically excludes all the severe lesions. Hence, they do not present a true picture. Our observation of prevalence of 26.4/1000 patients cannot be compared to earlier studies, because we included all mild, moderate and severe CHDs in age groups ranging from 0 to 15 years.

5. References

1. Ferencz C, Rubin JD, McCarter RJ, Brenner JI, Neil CA, Perry LW *et al.* Congenital heart disease: prevalence at live birth. The Baltimore-Washington Infant Study. *Am J Epidemiol* 2010; 121:31-36.
2. Bolisetty S, Daftary A, Ewald D, Knight B, Wheaton G. Congenital heart defects in Central Australia. *Med J Aust* 2014; 180:614-617.
3. Saxena A. Congenital heart disease in India: A status report. *Indian J Pediatr* 2012; 72:595-598.
4. Khalil A, Aggarwal R, Thirupuram S, Arora R. Incidence of congenital heart disease among hospital live births in India. *Indian Pediatr* 2014; 31:519-527.
5. Gupta I, Gupta ML, Parihar A, Gupta CD. Epidemiology of rheumatic and congenital heart disease in school children. *J Indian Medical Assoc* 2012; 90:57-59.
6. Vashishtha VM, Kalra A, Kalra K, Jain VK. Prevalence of congenital heart disease in school children. *Indian Pediatr* 2013; 30:1337-1340.
7. Jose VJ, Gomathi M. Declining prevalence of rheumatic heart disease in rural school children in India: 2011-2012. *Indian Heart J* 2013; 55:158-160.
8. Vaidyanathan B, Krishna kumar R. The global burden of congenital heart disease. *Congenital Cardiology Today* 2005; 3:1-8.