



Comparative study of long term consumption of *Corchorus olitorius* (Ewedu) and *Ocimum gratissimum* (Efirin) Diet-inclusion on male wistar rat

Kunle Ogungbemi¹, Ilesanmi FF², Odeniyi TA³, Ilori AO⁴, Oke OA⁵, Ogungbemi AM⁶, Balogun B⁷, Ogunremi OB⁸

^{1-5,7}Department of Chemistry/Biochemistry, Nigerian Stored Product Research Institute, Ibadan, Nigeria

⁶Department of Crop Production and Soil Science, Ladoko Akintola University of Technology, Ogbomosho, Oyo State, Nigeria

⁸Department of Outreach, Nigerian Stored Product Research Institute, Ibadan Oyo State, Nigeria

Abstract

This study was carried out to determine the effects of long-term consumption of different levels of *C. olitorius* and *O. gratissimum* diet-inclusion on male Wistar rat. A total of thirty-five (35) rats (average weight of 80 g±10g) were purchased and randomly allotted into seven (7) treatment groups with graded levels of *C. olitorius* and *O. gratissimum* (2.5%, 5%, and 10%) while the control had neither *C. olitorius* nor *O. gratissimum*. The feed intakes, weight performance as well as their histology of the experimental rats were monitored for eight weeks. The results obtained showed that the highest level of feed intake was recorded for animals in the group fed 10% level of *C. olitorius* supplementation when compared to other treatment groups. A similar result was observed for the weight performance which can be attributed to the efficient utilization of feed consumption. Also, group of rats fed with the lowest supplementation (2.5%) of *O. gratissimum* showed a significant increase ($P < 0.05$) in the feed intake than the group fed with the highest supplementation (10%). Also, the weight gain by the rats fed 2.5% *O. gratissimum* diet supplementation had the highest weight gain. The observable increase in the feed intake, weight gain and feed efficiency seen in the lowest supplementation level (2.5%) can be attributed to the spicing and seasoning of food with *O. gratissimum* and as such could also increase the palatability of the food. Therefore, it can be concluded that the leaves of *C. olitorius* are palatable and highly nutritious and can improve growth performance. Whereas, it can also be found that 2.5% *O. gratissimum* supplementation has better palatability than other experimental diets which reflected on the performance characteristics of this treatment and imposed an adverse effect with increasing dosage at 10% *O. gratissimum*. Thus, intake of *C. olitorius* should be encouraged while *O. gratissimum* is advised to be taken as a spice, not as a soup.

Keywords: *Corchorus olitorius*, *Ocimum gratissimum*, feed intake, feed efficiency, weight performance, wistar rat

Introduction

Plant foods especially vegetables contribute substantially to both local diets and ethno-medicine in developing countries especially Nigeria. (Okafor, 1980) ^[1]. Several vegetable have been used in traditional medicine worldwide and is now recognized by World Health Organization (WHO) as an essential building block for primary healthcare Onayade *et al.*, (1990) ^[2]. According to the WHO, more than 80% of the world population still relies on naturally occurring medicinal plants as their primary source of healthcare Adeyemi *et al.*, (2009) ^[3]. Thus, dietary plant plays an important role in healthcare management system. Dietary medicinal plant, thus are plants that possess some substances either in their leaves, stem, root etc. that can therefore be used in the treatment of ailments. Pharmacognostical evaluation of *O. gratissimum* used as traditional medicine in various diseases was drawn up according to WHO guidelines Matasyoh, (2012) ^[19]. Additionally, recent study on the phytochemical of this plant by Barku *et al.*, (2013) ^[18] shows the presence of polyphenols and flavonoid and other phytochemical constituent similarly to that of *C. olitorius* Lin *et al.*, (2002). The presence of these materials in these plants accounts for their usefulness as Medicinal plants. *C. olitorius* (Tiliaceae) is an annual herb whose leaves and roots are used as herbal medicine and eaten as vegetable by local people in East Malaysia, India, Egypt, and Philippines. (Zeghichi *et al.*,

2003) ^[4] In West African countries particularly Ghana, Nigeria and Sierra Leone, where staple diets consist of starchy food-stuffs such as rice, cassava, maize and yams, leafy vegetables are used to complement such staple foods (Tulio *et al.*, 2002) ^[5]. Traditionally, its leaves are used in the treatment of pain, fever, chronic cystitis and tumors (Abu-Hadid *et al.*, 1994) ^[6]. *C. olitorius* leaves are rich in antioxidants, fatty acids, minerals, vitamins and mucilaginous polysaccharides (Matsufuji *et al.*, 2001) ^[7]. They have become a part of daily diet improving the metabolic and physiological processes in the human body including in the weight performance. Similarly, in Africa *O. gratissimum* belongs to the family Lamiaceae and is widely distributed in tropical and warm regions of the world (Okigbo and Ogbonnaya, 2006) ^[8]. Among the numerous functions and role played by *C. olitorius*, the impact of this plant on weight performance and its palatability is yet to be assessed. Thus, it is expedient to assess the feed intake and weight performance in male wistar rats fed with different levels of *C. olitorius* diet inclusion. Also, pumpkin is often recommended by veterinarians as a dietary supplement for dogs and cats that are experiencing digestive problems. The high fiber content helps to aid proper digestion. Raw pumpkin can be fed to poultry as a supplement to regular feed, during the winter to help maintain egg production, which usually drops off during the cold months. Similarly, in Africa *O. gratissimum* belongs to the family Lamiaceae

and is widely distributed in tropical and warm regions of the world (Okigbo and Ogbonnaya, 2006) [8]. Leaf of *O. gratissimum* have been found to contain methylchavicol, linalool, eugenol, thymol and xanthamicrol and the amount produced is dependent on the area that it is cultivated as well as part of the plant (Okujagu *et al.*, 2008, Odebiyi and Sofowora, 1977) [9, 10], It is mainly used as a spice to flavor foods and meats. (Okigbo, 1975) [11] Their use as spice is known to reduce microbial load and extend the shelf life of foods. A lot of research has been done on *O. gratissimum* especially on its medicinal properties generating an abundance of information. However there is a need to see the effect of long term consumption of this plants on growth performance and its palatability via the intake

Methodology

Materials and Methods

Experimental Animals

A total number of twenty (35) Wistar rats weighing 70-90g were purchased and housed in the Faculty of Pharmaceutical Science Animal house, Ahmadu Bello University, Zaria, Nigeria and allowed free access to water and standard diet for three weeks for acclimatization. The study was conducted in accordance with the US guideline as contained in the National Institute of Health guide for the care and use of laboratory animals NIH publication No. 18-23, 1985. Chah *et al.*, (2006) [12]

Collection of Plant materials

The fresh plants were collected from a local farm in Agwan Jeba, Zaria, Kaduna State, Nigeria. Identification and authentication were carried out in the Herbarium Section of the Department of Botany, Ahmadu Bello University, Zaria, Nigeria, where a voucher number 2649 was assumed

Processing of Plant Materials

The fresh leaves of the *C. olitorius* and *O. gratissimum* were air-dried at room temperature 25°C. They were grounded into fine powder using an electric blender and stored in a dry sample container.

Formulation of experimental diets

Two experimental diets namely, the control diet (which contained 0% inclusion of *C. olitorius*, and *O. gratissimum*) *C. olitorius*, and *O. gratissimum* supplemented diet which was varied according to according to (Yan *et al.*, 1998) at 2.5%, 5%, and 10%.

Table 1: The cumulative body weight performance, feed intake and feed efficiency of wister rats fed with different levels of *C. olitorius* diet inclusion for 8weeks

Treatment	body weight gain (g/rat)	feed intake (g/rat)	feed efficiency (gain/food)
Control feed	41.60±3.60ab	109.09 ± 1.55a	0.3813±0.056a
2.5% <i>C. olitorius</i> diet Supplementation	43.47±3.77a	115.56 ± 1.56b	0.3760±0.048a
5% <i>C. olitorius</i> diet Supplementation	44.02±5.29ab	115.43 ± 4.92ab	0.3800±0.073a
10% <i>C. olitorius</i> diet Supplementation	56.05±6.30b	128.078± 5.54a	0.438±0.032b

Values with different superscript down the column are significantly different (P<0.05) with n=5

The body weight performance, feed intake and feed efficiency of wister rats fed with different levels of *O. gratissimum* diet inclusion as seen in Table 2. From the result obtained, a consistent increase was observed in the weight performance of animal fed with normal control (0%

Determination of Feed intake

a. The average of the feed intake (AFI) in the various supplemented groups and the control group for 8weeks was calculated, the average weight gain (AWG) of the animal in their respective group was also calculated. The Feed intake (g/kg body weight/week) was extrapolated as

$$\text{Feed intake} = \left(\frac{\text{Average feed intake (g)}}{\text{Average Body weight (kg) of rat}} \times 1000 \right) / 8\text{weeks}$$

Determination of Weight performance (weight change)

a. The weekly weight change was evaluated by subtracting the initial average weight of the animals (IAW) to the final average weight of the animal (FAW) on weekly basis

$$\text{Average weekly weight change} = \text{FAW} - \text{IAW}$$

b. The average cumulative weight change (ACWC) was evaluated by summing up the result obtained from (FAW-IAW)

Determination of Feed Efficiency

The weight changes by the animal in respectful groups were sum up on weekly basis to the end of the study (8 weeks). The weekly feed intakes were calculated for the supplemented and control groups and the feed efficiency is evaluated as:

$$\text{Feed efficiency} = \left(\frac{\text{weight gain (g/rat)}}{\text{Feed intake (g/rat)}} \right)$$

Statistical Analysis

The statistical significance between the control and other groups of experimental animals were determined by one-way analysis of variance (ANOVA) followed by Bonferroni t-test for multiple comparisons. The results are presented as mean ± SD at confidence level of 95% (p≤0.05)

Results

The cumulative body weight performance, feed intake and feed efficiency of wister rats fed with different levels of *C. olitorius* diet inclusion as seen in Table 1. The result obtained showed group fed with the highest feed intake gave the highest weight performance when compared with animal fed with Normal control (0% *C. olitorius* inclusion) and 2.5% and 5% *C. olitorius* diet inclusion and it turns, the feed efficiency increases with increasing diet inclusion.

O. gratissimum) and 2.5% *O. gratissimum* diet inclusion while the highest *O. gratissimum* diet inclusion (10%) showed a decrease in its weight performance as the supplementation tally to 8 weeks when compared to groups fed with lowest *O. gratissimum* diet inclusion.

Table 2: The cumulative weight performance, feed intake (g/rat) and feed efficiency of wister rats fed with different levels of *O. gratissimum* diet inclusion for 8weeks

Treatment	body weight gal (g/rat)	feed intake (g/rat)	feed efficiency (gain/food)
Control feed	41.60±3.60ab	109.09 ± 1.59	0.3813±0.056a
2.5% <i>O. gratissimum</i> diet Supplementation	47.32±4.99a	120.09 ± 1.56b	0.3940±0.039
5% <i>O. gratissimum</i> diet Supplementation	44.77±1.77a	115.29 ± 4.92ab	0.3893±0.042a
10% <i>O. gratissimum</i> diet Supplementation	33.07±6.20b	108.50 ± 5.54a	0.3047±0.051b

Values with different superscript down the column are significantly different ($P < 0.05$) with $n=5$

Discussion

The utilization of leaves of *C. olitorius* and *O. gratissimum* added to a standard feed (vital feed growers mash) by male wister rats gave some substantial results on the feed intake, feed efficiency, weight gain, as seen in results obtained from this findings. There was an appreciable increase in the feed intake, weight performance and feed efficiency obtained in rats fed with the *C. olitorius* diet supplementation with 10% *C. olitorius* having the highest feed rate which can be attributed to leaves of *C. olitorius* that possesses good amount of minerals and vitamins Ndlovu *et al.*, (2008) [14] and Shitanda *et al.*, (2006) [15], which plays a role of stimulating the appetite of the rats. Also, *C. olitorius* plays a supplementary role adding variety to the diets such as good flavor and improve palatability and taste of the main staple food hence broadening the feed intake Iyimo *et al.*, (2003) [16]. This dietary plant plays an important role in healthcare management system and substantially, in the weight performance and feed efficiency due to its palatability and its nutritional endowments. Thus, it can be concluded that plant *C. olitorius* enhances weight gain and improves the look of the staple food taken in return improve feed efficiency. Whereas, The long term consumption of *O. gratissimum* diet inclusion by male Wister rats had some considerable effect on the weight/growth performance and also on the rate of their feed intake as seen in the results obtained from the studies. It was observed that *O. gratissimum* diet inclusion with vital feed resulted in increase weight gain seen in Table 2 which is attributed to the nutrient composition of the leaves of *O. gratissimum*. *O. gratissimum* contain nutrient that can increase weight such as carbohydrate, protein and lipid, minerals and vitamin Edeoga *et al.*, (2006). The increase in the feed intake observed in Table 2 could be explained from the fact that *O. gratissimum* leaves are rich in minerals and vitamins Adepoju and Oyewole, (2008) [17], which also stimulated the appetite of the rats. Ibrahim *et al.*, (1997) reported that *O. gratissimum* could be used for spicing and seasoning of food and as such could also increase the palatability of food which is in agreement with the result gotten showing 2.5% supplemented diet group having the highest intake. The aroma from the *O. gratissimum*, which are mainly eugenol methyl cinnamate, camphor and thymol Nakamura *et al.*, (1999), might have some feed digestive properties that increase the appetite of the rats but at a very high dosage, the diet can become repulsive which probably made 2.5% supplementation the best giving a growth performance resulting from the palatability potentials of the spice (2.5%) of *O. gratissimum* Inclusion.

Histology result as seen in plate (1-4) shows that *C. olitorius* does not in any way affect the architecture of the colon whereas *O. gratissimum* at 10% shows an adverse effect on the architecture of the colon with some abnormal cell

absorbing maximally the die to give different color to other cells in the normal control.

References

- Okafor JC. Edible indigenous woody plants in the rural economy of the Nigerian forest zone. *Forest Ecol Manag.* 1980; 3:45-55.
- Onayade OA, Scheffer JJC, Baerheim Svendsen A. Importance of phytotherapy and screening of plants used medicinally in Africa. *Planta Medica.* 1990; 56(6):doi:10.1055/s-002-5459.
- Adeyemi OS, Akanji MA, Oguntoye SA. Ethanolic leaf extract of *Psidium guajava*: Phytochemical and trypanocidal activity in rats infected with *Trypanosoma Brucei Brucei*. *Journal of medicinal plants research.* 2009; 3(5):420-3.
- Zeghichi S, Kallithraka S, Simopoulos AP. Nutritional composition of molokhia (*C. olitorius*) and stamnagathi (*Cichorium spinosum*). *Plants in Human Health and Nutrition policy.* *World Rev Nutr Diet.* 2003; 91:1-21.
- Tulio AZ, Ose K, Chachin K, Ueda Y. Effects of storage temperatures on the postharvest quality of jute leaves (*Corchorus olitorius* L.). *Postharvest Biol Technol.* 2002; 26(3):329-38.
- Abu-Hadid AF, El-Shinawy MZ, El-Bethagy AS, Gaafer SA, Medany M. Studies on the production of off-season Jew's mallow (molokhia) in Egypt. *Egyptian J Hort.* 1994; 21:187-193.
- Matsufuji H, Sakai S, Chino M, Goda Y, Toyoda M, Takeda M. Relationship Between Cardiac Glycoside Contents and Color of *Corchorus olitorius* Seeds. *J Health Sci.* 2001; 47(2):89-93.
- Okigbo RN, Ogbonnaya UO. Antifungal effects of two tropical plant leaf extracts (*O. gratissimum* and *Aframomum melegueta*) on postharvest yam (*Dioscorea* spp.) rot. *African J Biotechnol.* 2006; 5(9):727-31.
- Okujagu TF, Etaturvie SO, Eze I, Iimoh B, Nwokereke C, Mbaoji C, *et al.* *Medicinal Plants of Nigeria (South West Zone Vol. 1).* Nigeria Natural Medicine Development Agency, Federal Ministry of Science and Technology, Lagos, Nigeria, 2008, 204.
- Odebiyi OO, Sofowora EA. Phytochemical screening of Nigerian medicinal plants II. *Lloydia.* 1977; 41(3):234-46.
- Okigbo BN. Neglected plants of horticultural and nutritional importance in traditional farming systems of tropical Africa. In *IV Africa Symposium on Horticultural Crops.* 1975; 53:131-50.
- Chah KF, Eze CA, Emuelosi CE, Esimone CO. Antibacterial and wound healing properties of methanolic extracts of some Nigerian medicinal plants. *J Ethnopharmacol.* 2006; 104(1):164-7.

13. Yan L, Yee JA, Li D, McGuire MH, Thompson LU. Dietary flaxseed supplementation and experimental metastasis of melanoma cells in mice. *Cancer Letters*. 1998; 124(2):181-6.
14. Ndlovu J, Afolayan AJ. Nutritional analysis of the South African wild vegetable *C. olerius* L. *Asian J Plant Sci*. 2008; 7(6):615-8.
15. Shitanda D, Wanjala NV. Effect of different drying methods on the quality of jute (*C. olerius* L.). *Drying Technol*. 2006; 24(1):95-8.
16. Lyimo M, Temu RPC, Mugula JK. Identification and nutrient composition of indigenous vegetables of Tanzania. *Plant Foods Human Nutr*. 2003; 58(1):85-92.
17. Adepoju OT, Oyewole EO. Nutritional importance and Micronutrients potentials of non-conventional indigenous green leafy vegetable from Nigeria. *Agricultural J*. 2008; 3(5):362-5.
18. Barku VY, Boye A, Ayaba S. Phytochemical screening and assessment of wound healing activity of the leaves of *Anogeissus leiocarpus*. *European Journal of Experimental Biology*. 2013; 3(4):18-25.
19. Matasyoh LG. Genetic variation and medicinal activity in *O. gratissimum* l. of kenya (Doctoral dissertation), 2012.
20. Li CJ, Huang SY, Wu MY, Chen YC, Tsang SF, Chyuan JH, *et al*. Induction of apoptosis by ethanolic extract of *C. olerius* leaf in human hepatocellular carcinoma (HepG2) cells via a mitochondria-dependent pathway. *Molecules*. 2012; 17(8):9348-9360.