

On the negative Pell equation $y^2 = 7x^2 - 14$

¹ MA Gopalan, ² KK Viswanathan, ³ P Gayathri

¹ Professor, Department of Mathematics, SIGC, Trichy, Tamil Nadu, India

² Associate Professor, Kuwait College of Science and Technology, Safat, Kuwait

³ M.Phil Scholar, Department of Mathematics, SIGC, Trichy, Tamil Nadu, India

Abstract

The binary quadratic equation represented by the negative pellian $y^2 = 7x^2 - 14$ is analyzed for its distinct integer solutions. A few interesting relations among the solutions are also given. Further, employing the solutions of the above hyperbola, we have obtained solutions of other choices of hyperbolas, parabolas and special Pythagorean triangle.

Keywords: binary quadratic, hyperbola, parabola, integral solutions, pell equation

Introduction

Diophantine equation of the form $y^2 = Dx^2 + 1$, where D is a given positive square - free integer is known as pell equation and is one of the oldest Diophantine equation that has interesting mathematicians all over the world, since antiquity, J.L.Lagrange proved that the positive pell equation $y^2 = Dx^2 + 1$ has infinitely many distinct integer solutions whereas the negative pell equation $y^2 = Dx^2 - 1$ does not always have a solution. In ^[1], an elementary proof of a criterium for the solvability of the pell equation $x^2 - Dy^2 = -1$ where D is any positive non-square integer has been presented. For examples the equations $y^2 = 3x^2 - 1$, $y^2 = 7x^2 - 4$ have no integer solutions whereas $y^2 = 65x^2 - 1$, $y^2 = 202x^2 - 1$ have integer solutions. In this context, one may refer ^[2-17]. More specifically, one may refer "The On-line Encyclopedia of integer sequences" (A031396, A130226, A031398) for values of D for which the negative pell equation $y^2 = Dx^2 - 1$ is solvable or not.

In this communication, the negative Pell equation given by $y^2 = 7x^2 - 14$ is considered and infinitely many integer solutions are obtained. A few interesting relations among the solutions are presented.

Method of Analysis

The negative pell equation representing hyperbola under consideration is

$$y^2 = 7x^2 - 14 \quad (1)$$

whose smallest positive integer solution is $x_0 = 3, y_0 = 7$

To obtain the other solutions of (1), consider the pell equation $y^2 = 7x^2 + 1$ whose solution is given by

$$\tilde{x}^n = \frac{g_n}{2\sqrt{7}}, \tilde{y}^n = \frac{f_n}{2}$$

Where,

$$f_n = (8 + 3\sqrt{7})^{n+1} + (8 - 3\sqrt{7})^{n+1}$$

$$g_n = (8 + 3\sqrt{7})^{n+1} - (8 - 3\sqrt{7})^{n+1}$$

Applying Brahamagupta Lemma between (x_0, y_0) and $(\tilde{x}^n, \tilde{y}^n)$, the other integer solutions of (1) are given by

$$x_{n+1} = \frac{3}{2}f_n + \frac{\sqrt{7}}{2}g_n,$$

$$y_{n+1} = \frac{7}{2}f_n + \frac{3\sqrt{7}}{2}g_n, n = 0, 1, 2, \dots$$

The recurrence relations satisfied by the solutions x and y are given by

$$x_{n+3} - 16x_{n+2} + x_{n+1} = 0$$

$$y_{n+3} - 16y_{n+2} + y_{n+1} = 0$$

Some numerical examples of x and y satisfying (1) are given in the Table1 below.

Table 1: Examples

n	x_n	y_n
0	3	7
1	45	119
2	717	1897
3	11427	30233
4	182115	481831

From the above table, we observe some interesting relations among the solutions which are presented below.

- Both x_n & y_n values are always odd.
- Each of the following expressions is a nasty number:

- $2[17x_{2n+2} - x_{2n+3}] + 12$
- $\frac{271x_{2n+2} - x_{2n+4} + 96}{8}$
- $\frac{135x_{2n+2} - 3y_{2n+3} + 48}{4}$
- $\frac{4302x_{2n+2} - 6y_{2n+4} + 1524}{127}$
- $(542x_{2n+3} - 34x_{2n+4}) + 12$
- $\frac{9x_{2n+3} - 51y_{2n+2} + 48}{4}$
- $270x_{2n+3} - 102y_{2n+3} + 12$
- $\frac{2151x_{2n+3} - 51y_{2n+4} + 48}{4}$
- $\frac{18x_{2n+4} - 1626y_{2n+2} + 1524}{127}$
- $\frac{135x_{2n+4} - 813y_{2n+3} + 48}{4}$
- $4302x_{2n+4} - 1626y_{2n+4} + 12$
- $\frac{6y_{2n+3} - 90y_{2n+2} + 84}{7}$
- $\frac{3y_{2n+4} - 717y_{2n+2} + 672}{56}$
- $\frac{90y_{2n+4} - 1434y_{2n+3} + 84}{7}$

- Each of the following expression is a cubical integer:

- $9(17x_{3n+3} - x_{3n+4}) + 9(17x_{n+1} - x_{n+2})$
- $2304(271x_{3n+3} - x_{3n+5}) + 6912(271x_{n+1} - x_{n+3})$
- $(45x_{3n+3} - y_{3n+4}) + 3(45x_{n+1} - y_{n+3})$

- $16129(717x_{3n+3} - y_{3n+5}) + 48387(717x_{n+1} - y_{n+3})$
- $9(271x_{3n+4} - 17y_{3n+5}) + 27(271x_{n+2} - 17y_{n+3})$
- $(3x_{3n+4} - 17y_{3n+3}) + 3(3x_{n+2} - 17y_{n+1})$
- $(45x_{3n+4} - 17y_{3n+4}) + 3(45x_{n+2} - 17y_{n+2})$
- $(717x_{3n+4} - 17y_{3n+5}) + 3(717x_{n+2} - 17y_{n+3})$
- $16129(3x_{3n+5} - 271y_{3n+3}) + 48387(3x_{n+3} - 271y_{n+1})$
- $(45x_{3n+5} - 271y_{3n+4}) + 3(45x_{n+3} - 271y_{n+2})$
- $(717x_{3n+5} - 271y_{3n+5}) + 3(717x_{n+3} - 271y_{n+3})$
- $49(y_{3n+4} - 15y_{3n+3}) + 147(y_{n+2} - 15y_{n+1})$
- $12544(y_{3n+5} - 239y_{3n+3}) + 37632(y_{n+3} - 239y_{n+1})$
- $441(45y_{3n+5} - 717y_{3n+4}) + 1323(45y_{n+3} - 717y_{n+2})$

4. Relations among the solutions:

- $2x_{n+3} = 32x_{n+2} - 2x_{n+1}$
- $6y_{n+1} = 2x_{n+2} - 16x_{n+1}$
- $6y_{n+2} = 16x_{n+2} - 2x_{n+1}$
- $6y_{n+3} = 254x_{n+2} - 16x_{n+1}$
- $96x_{n+2} = 6x_{n+1} + 6x_{n+3}$
- $96y_{n+1} = 2x_{n+3} - 254x_{n+1}$
- $96y_{n+2} = 16x_{n+3} - 16x_{n+1}$
- $96y_{n+3} = 254x_{n+3} - 2x_{n+1}$
- $16x_{n+2} = 2x_{n+1} + 6y_{n+2}$
- $16y_{n+1} = 2y_{n+2} - 42x_{n+1}$
- $16y_{n+3} = 42x_{n+1} + 254y_{n+2}$
- $254y_{n+1} = 2y_{n+3} - 672x_{n+1}$
- $6y_{n+1} = 16x_{n+3} - 254x_{n+2}$
- $6y_{n+2} = 2x_{n+3} - 16x_{n+2}$
- $6y_{n+3} = 16x_{n+3} - 2x_{n+2}$
- $16y_{n+2} = 42x_{n+2} + 2y_{n+1}$
- $16y_{n+3} = 672x_{n+2} + 16y_{n+1}$
- $2y_{n+3} = 42x_{n+2} + 16y_{n+2}$
- $254y_{n+2} = 42x_{n+3} + 16y_{n+1}$
- $254y_{n+3} = 672x_{n+3} + 2y_{n+1}$
- $16y_{n+3} = 42x_{n+3} + 2y_{n+2}$
- $42y_{n+3} = 672y_{n+2} - 42y_{n+1}$
- $10752x_{n+1} = 32y_{n+3} - 4064y_{n+1}$
- $10752x_{n+2} = 256y_{n+3} - 256y_{n+1}$
- $10752x_{n+3} = 4064y_{n+3} - 32y_{n+1}$

- $10752y_{n+2} = 672y_{n+3} + 672y_{n+1}$
- $42y_{n+1} = 672y_{n+2} - 42y_{n+3}$

Remarkable Observations

i) Employing linear combinations among the solutions of (1), one may generate integer solutions for other choices of hyperbolas which are presented in the Table2 below.

Table 2: Hyperbolas

S. No.	(X _n , Y _n)	Hyperbola
1	$(x_{n+2} - 15x_{n+1}, 17x_{n+1} - x_{n+2})$	$7Y^2 - 9X^2 = 252$
2	$(x_{n+3} - 239x_{n+1}, 271x_{n+1} - x_{n+3})$	$1792Y^2 - 2304X^2 = 16515072$
3	$(3y_{n+2} - 119x_{n+1}, 45x_{n+1} - y_{n+2})$	$448Y^2 - 64X^2 = 114688$
4	$(3y_{n+3} - 1897x_{n+1}, 717x_{n+1} - y_{n+3})$	$112903Y^2 - 16129X^2 = 7284049948$
5	$(45x_{n+3} - 717x_{n+2}, 271x_{n+2} - 17x_{n+3})$	$63Y^2 - 9X^2 = 2268$
6	$(45y_{n+1} - 7x_{n+2}, 3x_{n+2} - 17y_{n+1})$	$448Y^2 - 64X^2 = 28672$
7	$(45y_{n+2} - 119x_{n+2}, 45x_{n+2} - 7y_{n+2})$	$7Y^2 - X^2 = 28$
8	$(45y_{n+3} - 1897x_{n+2}, 717x_{n+2} - 17y_{n+3})$	$448Y^2 - 64X^2 = 114688$
9	$(717y_{n+1} - 7x_{n+3}, 3x_{n+3} - 271y_{n+1})$	$112903Y^2 - 16129X^2 = 7284049948$
10	$(717y_{n+2} - 119x_{n+3}, 45x_{n+3} - 271y_{n+2})$	$448Y^2 - 64X^2 = 114688$
11	$(717y_{n+3} - 1897x_{n+3}, 717x_{n+3} - 271y_{n+3})$	$7Y^2 - X^2 = 28$
12	$(17y_{n+1} - y_{n+2}, y_{n+2} - 15y_{n+1})$	$21Y^2 - 49X^2 = 4116$
13	$(271y_{n+1} - y_{n+3}, y_{n+3} - 239y_{n+1})$	$16128Y^2 - 12544X^2 = 809238528$
14	$(1897y_{n+2} - 119y_{n+3}, 45y_{n+3} - 717y_{n+2})$	$7Y^2 - X^2 = 12348$

ii) Employing linear combinations among the solutions of (1), one may generate integer solutions for other choices of parabolas which are presented in the Table3 below.

Table 3: Parabolas

S. No.	(X _n , Y _n)	Parabola
1	$(x_{n+2} - 15x_{n+1}, 17x_{2n+2} - x_{2n+3})$	$3X^2 = 7Y - 42$
2	$(x_{n+3} - 239x_{n+1}, 271x_{2n+2} - x_{2n+4})$	$48X^2 = 1792Y - 172032$
3	$(3y_{n+2} - 119x_{n+1}, 45x_{2n+2} - y_{2n+3})$	$8X^2 = 448Y - 7168$
4	$(3y_{n+3} - 1897x_{n+1}, 717x_{2n+2} - y_{2n+4})$	$127X^2 = 112903Y - 28677362$
5	$(45x_{n+3} - 717x_{n+2}, 271x_{2n+3} - 17x_{2n+4})$	$3X^2 = 63Y - 378$
6	$(45y_{n+1} - 7x_{n+2}, 3x_{2n+3} - 17y_{2n+2})$	$8X^2 = 448Y - 7168$
7	$(45y_{n+2} - 119x_{n+2}, 45x_{2n+3} - 17y_{2n+3})$	$X^2 = 7Y - 14$
8	$(45y_{n+3} - 1897x_{n+2}, 717x_{2n+3} - 17y_{2n+4})$	$8X^2 = 448Y - 7168$
9	$(717y_{n+1} - 7x_{n+3}, 3x_{2n+4} - 271y_{2n+2})$	$127X^2 = 112903Y - 28677362$
10	$(717y_{n+2} - 119x_{n+3}, 45x_{2n+4} - 271y_{2n+3})$	$8X^2 = 448Y - 7168$
11	$(717y_{n+3} - 1897x_{n+3}, 717x_{2n+4} - 271y_{2n+4})$	$X^2 = 7Y - 14$
12	$(17y_{n+1} - y_{n+2}, y_{2n+3} - 15y_{2n+2})$	$7X^2 = 21Y - 294$
13	$(271y_{n+1} - y_{n+3}, y_{2n+4} - 239y_{2n+2})$	$112X^2 = 16128Y - 3612672$
14	$(1897y_{n+2} - 119y_{n+3}, 45y_{2n+4} - 717y_{2n+3})$	$X^2 = 147Y - 6174$

iii) Consider $m = x_{n+1} + y_{n+1}$, $n = x_{n+1}$, observe that $m > n > 0$. Treat m, n as the generators of the Pythagorean triangle

$$T(\alpha, \beta, \gamma), \text{ where } \alpha = 2mn, \beta = m^2 - n^2, \gamma = m^2 + n^2$$

Then the following interesting relations are observed.

a) $2\alpha - 7\beta + 5\gamma = 28$

b) $9\alpha - 2\gamma - \frac{28A}{P} = 28$

c) $\frac{2A}{P} = x_{n+1}y_{n+1}$

Conclusion

In this paper, we have presented infinitely many integer solutions for the hyperbola represented by the negative Pell equation $y^2 = 7x^2 - 14$. As the binary quadratic Diophantine equation are rich in variety, one may search for the other choices of negative Pell equations and determine their integer solutions along with suitable properties.

References

1. Mollin RA, Anitha Srinivasan. A note on the Negative Pell Equation, International Journal of Algebra. 2010; 4(19):919-922.
2. Whitford EE. Some Solutions of the Pellian Equations $x^2 - Ay^2 = \pm 4$ JSTOR: Annals of Mathematics, Second Series. 1913-1914; 1:157-160.
3. Ahmet Tekcan S. Betw Gezer and Osman Bizim. On the Integer Solutions of the Pell Equation $x^2 - dy^2 = 2^t$, World Academy of Science, Engineering and Technology. 2007; 1:522-526.
4. Ahmet Tekcan. The Pell Equation $x^2 - (k^2 - k)y^2 = 2^t$. World Academy of Science, Engineering and Technology. 2008; 19:697-701.
5. Merve Guney. Solutions of the Pell Equation $x^2 - (a^2b^2 + 2b)y^2 = 2^t$, when $N \in (\pm 1, \pm 4)$, Mathematics Aterna. 2012; 2(7):629-638.
6. Sangeetha v, Gopalan MA, Manju Somanath. On the Integral Solutions of the Pell Equation $x^2 = 13y^2 - 3^t$, International Journal of Applied Mathematics Research. 2014; 3(1):58-61.
7. Gopalan MA, Sumathi G, Vidhyalakshmi S. Observations on the Hyperbola $x^2 = 19y^2 - 3^t$, Scholars Journal of the Engineering and Technology. 2014; 2(2A):152-155.
8. Gopalan MA, Vidhyalakshmi S, Kavitha A. On the Integral Solution of the Binary Quadratic Equation $x^2 = 15y^2 - 11^t$, Scholars Journal of the Engineering and Technology. 2014; 2(2A):156-158.
9. Vidhyalakshmi S, Karthiga V, Agalya K. On the Negative Pell Equation, Proceedings of the National Conference on MATAM. 2015, 4-9.
10. Meena K, Gopalan MA, Karthika R. On the Negative Pell Equation $y^2 = 10x^2 - 6^t$, International Journal of Multidisciplinary Research and Development. 2015; 2(12):390-392.
11. Gopalan MA, Vidhyalakshmi S, Pandichelvi V, Sivakamasundari P, Priyadharsini C. On the Negative Pell Equation $y^2 = 45x^2 - 11^t$, International Journal of Pure Mathematical Science. 2016; 16:30-36.
12. Meena K, Vidhyalakshmi S, Rukmani A. On the Negative Pell Equation $y^2 = 31x^2 - 6^t$, Universe of Emerging Technologies and Science. 2015; 2(12):1-4.
13. Meena K, Gopalan MA, Bhuvaneswari E. On the Negative Pell Equation $y^2 = 60x^2 - 15^t$, Scholars Bulletin. 2015; 1(11):310-316.
14. Gopalan MA, Vidhyalakshmi S, Presenna R, Vanitha M. Observations on the Negative Pell Equation $y^2 = 180x^2 - 11^t$, Universal Journal of Mathematics. 2(1):41-45.
15. Vidhyalakshmi S, Gopalan MA, Premalatha E, Sofiya christinal S. On the Negative Pell Equation $y^2 = 72x^2 - 8^t$, International Journal of Emerging Technologies in Engineering Research (IJETER). 2016; 4(2):25-28.
16. Gopalan MA, Vidhyalakshmi S, Thiruniraiselvi N. A study on the Hyperbola $y^2 = 8x^2 - 31^t$, International Journal of Latest Research in Science and Technology. 2013; 2(1):454-456.
17. Gopalan MA, Vidhyalakshmi S, Usha Rani TR, Mallika S. Observations on the Hyperbola $y^2 = 12x^2 - 3^t$, Besse Journal of Math. 2012; 2(3):153-158.